

Proximate Compositions and Biological Activities of *Caulerpa lentillifera***Risa Nofiani*¹, Sigit Hertanto¹, Titin Anita Zaharah¹, and Sutarman Gafur²**¹Department of Chemistry, Faculty of Mathematics and Natural Sciences,
Universitas Tanjungpura, Pontianak, Indonesia²Department of Soil Science, Faculty of Agriculture, Universitas Tanjungpura,
Pontianak, Indonesia

*email: risa.nofiani@chemistry.untan.ac.id, rnofiani@yahoo.com

Received September 14, 2018; **Accepted** November 25, 2018; **Available online** December 8, 2018**ABSTRACT**

Caulerpa lentillifera is an edible and functional seaweed due to its high nutritional compositions and its biological activities. In this study, *C. lentillifera* was evaluated for its proximate compositions (moisture, ash, protein, lipid and fiber contents) and its biological activities (antimicrobial, anti-oxidant, and toxicity). Moisture content, crude lipid, crude protein, and crude fiber were determined using oven method, soxhlet extraction, semi-micro Kjeldhal, and hydrolysis, respectively. Fresh *C. lentillifera* of Natuna Island, Indonesia, showed its higher level content of ash, crude lipid, and crude fiber compared to that of fresh *C. lentillifera* of Penghu, Taiwan. For its biological activity assays, the extracts were prepared from fresh and dry *C. lentillifera* (FC and DC). Both of the extracts showed the broad spectrum of weak antimicrobial using well-diffusion agar tests and antioxidant activities using a modified linoleic acid emulsion system. The toxicity for both extracts was determined using brine shrimp lethality test. DC extract showed its very low toxicity level and there was no toxicity for FC. Hemolytic activity was determined using red blood assay. Both extracts showed their low hemolytic activities (about 5-13%) for the concentration of 100 and 150 µg/mL, but the activity increased sharply (about 96%) on the concentration of 200 µg/mL. It was concluded that *C. lentillifera* has a potency as a functional food due to containing secondary metabolites with various biological activities.

Keywords: *Caulerpa lentillifera*, antimicrobial and antioxidant activity, toxicity, hemolysis, proximate composition

INTRODUCTION

Seaweeds have high nutritional constituents consisting of essential minerals, fatty acids, dietary fibers, amino acids and vitamins (Bhuiyan, Qureshi, Kamal, Aftab, & Siddique, 2016). Studies about chemical composition and nutritive value of edible seaweeds have increased due to their potential for food uses. For example, *C. lentillifera* from Malaysia and Thailand were reported having the nutritional composition such as: proximate composition, amino acids, vitamin C and E (Matanjan et al, 2009; Ratana-arporn and Chirapart, 2006). It was also reported that dry *C. lentillifera* (DC) which can be used as sources of food protein due to its high protein levels and balanced amino acid profiles (Ratana-arporn & Chirapart, 2006)

C. lentillifera is a green edible seaweed, classified into a Chlorophyta marine macroalgae (Mohamed, Hashim, & Rahman, 2012). It is widespread in Asian countries such as: Indonesia, Japan, Thailand, Malaysia, China, Philiphine, Korea and some other countries in Southeast Asia (Nguyen, Ueng, & Tsai, 2011).

In Natuna Island, Indonesia, *C. lentillifera* is naturally grown and known as Latuh and it is used to be consumed as a fresh salad, although it is lately rarely consumed by the community of Natuna.

In addition to its good nutritional composition, the genus *Caulerpa* is known rich in its bioactive compounds with various biological activities. These properties make it be potentially used as a functional food. *Caulerpa racemose* has anti-microbial activities due to its bioactive compounds content known as: caulerpin, sterol, and caulerpicin (Doty and Aguilar-santos, 1970; Doty 1966). *Caulerpa ashmeadii* contains high concentrations of sesquiterpenoid showing their anti-microbial and ichthyotoxicity activities (Paul, Littler, Littler, & Fenical, 1987). *Caulerpa taxipolia* and the other *Caulerpa* containing *caulerpenyne* showed their cytotoxic activities (Dumay, Erard, Pergent-martini, & Amade, 2002). *C. lentillifera* combined with *Hibiscus rosa-sinensis*, and *Piper sarmentosum* showed their antipyretic activities (Daud, Fazuliana, Arsad,

Ismail, & Tawang, 2016). Extract of *C. lentillifera* has the potency to be used as an anti-diabetic agent.

A seaweed grown in different location, habitat, or environmental conditions, and harvested in different maturity levels had caused the different in its chemical compositions, consistency, color, quality, and its bioactive compounds (Ito and Hori 2009; Kılınç et al. 2013). The proximate composition of *C. lentillifera* of Malaysia and Thailand has been reported but *C. lentillifera* of Indonesia has not yet been studied and reported.

In this study, *C. lentillifera* from Natuna Island of Indonesia was evaluated for its proximate compositions as *C. lentillifera* fresh because it is eaten fresh in the community of Natuna. For its biological activities particularly for its antimicrobial, anti-oxidant, and its toxicity activities was also analyzed as fresh and dry *C. lentillifera* which probably benefit for health or is called as a functional food.

EXPERIMENTAL SECTION

Sampling

C. lentillifera was collected randomly from the coastal area of Kabupaten Natuna, Kepulauan Riau. Morphology of *C. lentillifera* was identified in the Laboratory of Biology, Faculty of Mathematics and Natural Sciences, Universitas Tanjungpura, Indonesia.

Preparation of *C. lentillifera*

Fresh *C. lentillifera* (FC) was rinsed using distilled water and used directly for its proximate analysis and biological activity assays. Dry *C. lentillifera* (DC) were prepared by drying of *C. lentillifera* directly under the sunshine. The FC was used for proximate analysis, while the FC and DC were used for its antioxidant, hemolysis, cytotoxicity and antimicrobial assays.

Proximate Analysis

Moisture content was determined using oven method based on Method of 925.10 from AOAC. Ash content was determined based on Method number of 08-01 from AOAC. Crude lipid was determined using Soxhlet extraction with diethyl ether referred to Method of 30-25, AOAC. Crude fiber was determined using successive hydrolysis with 100°C 1.25% of H₂SO₄ and 3.25% of NaOH for 30 min each based on SNI 01-2891-1992. Crude protein was determined using semi-micro Kjeldhal

methods referred to SNI (Indonesian National Standard 01-2891-1992).

Extraction of *C. lentillifera*

The FC (139.6 g) was ground using a mortar and macerated using 80% of ethyl acetate. The maceration was carried out many times until colorless filtrate. The filtrate was evaporated using a rotary evaporator to get crude extract FC. The DC powder (21.6 g) was macerated using of methanol 80% until colorless filtrate. The filtrate was evaporated using a rotary evaporator to get crude extract DC.

Antimicrobial Test

Both extracts (FC and DC) were screened antimicrobial activities using well-diffusion methods (Okeke, Iroegbu, Eze, Okoli, & Esimone, 2001). Tested microorganisms were cultured by inoculation onto 10 mL of nutrient broth (NB) medium and incubated on a rotary shaker incubator at 130 rpm at 30 °C for 14-16 h. The culture was added on nutrient agar (NA), mixed and poured into the plate. After gel solid, the medium was punched using a sterilized pipette with diameter 0.4 mm. The extracts (FC and DC) with various concentrations (50, 100, 200, 400, and 500 µg/well) were poured into the well. After solvent from the extract was evaporated, the plate was incubated for 14-16 hours.

The extract having antimicrobial activities was signed with the formation of inhibition zone around the well then the diameter of inhibition zone (measured from the edge of the colony to the edge of the clear zone) was recorded. Furthermore, the extract having antimicrobial activities was tested for its bacteriostatic and bactericidal activities by the measuring its inhibition zone. Minimum inhibition concentrations (MICs) of each extract was counted based on the method of Bonev et al. (Bonev, Hooper, & Parisot, 2008).

Antioxidant Assay

The antioxidant assay was carried out using a slight modified linoleic acid emulsion system (Loganayaki, Siddhuraju, & Manian, 2013). A 10 µL of linoleic acid was mixed with 1 mL of ethanol in Eppendorf tube. The mixture was added with 1000 µg extract of *C. lentillifera* and incubated in the dark room at 25 °C.

$$\% \text{ of AA} = 1 - \frac{(\text{Sample absorbance at 24h} - \text{sample absorbance at 0 h})}{(\text{Blank absorbance at 24h} - \text{blank absorbance at 0 h})} \times 100\% \quad (1)$$

After 24 hours, the mixture was added with 20 μL of FeSO_4 0.014 M and 20 μL of KSCN. After 3 min, the absorbance was measured at 490 nm. A blank absorbance was measured at $t = 0$ h and $t = 24$ h. Vitamin C (1000 μg) was used as a positive control. The antioxidant activity (AA) was calculated as a percentage of inhibition relative to the control using the **Equation 1**.

Erythrocyte suspensions were prepared according to Situ & Bobek (2000) and Saengkhae et al. (2007). Cow blood was collected using a sterile container and added with an equal volume of Alsever's solution. The blood was centrifuged at $1.500 \times g$ for 5 menit with a temperature of 4 $^\circ\text{C}$. The pellet was washed three times with 10 volumes of phosphate buffer saline (PBS; 125 mM of NaCl, 10 mM of sodium phosphate buffer, pH 7.4). The pellet was resuspended in PBS solution and adjusted to a hematocrit of 1%.

Hemolytic Assay

The hemolytic assay was carried out using a red blood cell (Situ & Bobek, 2000; Saengkhae et al. 2007). A 100 μL of erythrocytes 1% was mixed with the extracts of FC and DC (100, 150 dan 200 $\mu\text{g}/\text{mL}$) and incubated at 37 $^\circ\text{C}$. After 30 min, the mixture was centrifuged at $1500 \times g$, 4 $^\circ\text{C}$. Cell lysis was monitored by measuring the release of hemoglobin at 540 nm using a Microplate reader. As a positive and negative control was used 1% of sodium dodecyl sulfate (SDS) in PBS and PBS alone, respectively. Percentage of hemolysis was calculated as follows: $[(A_{540} \text{ of the sample treated with the extract} - A_{540} \text{ of the sample treated with the extract}) / [(A_{540} \text{ of the sample treated with SDS} - A_{540} \text{ sample treated with buffer})] \times 100\%$.

sample treated with buffer)] / [(A_{540} of the sample treated with SDS - A_{540} sample treated with buffer)] x 100%.

Preliminary Toxicity Test

Preliminary toxicity test was carried out using Brine Shrimp Lethality Test (BSLT) (Meyer et al., 1982). Ten shrimps *Artemia salina* Leach, of 2 days old, were added with the extracts of various concentrations (0 (for control), 10, 100, 200, 400, 600, 800 dan 1.000 $\mu\text{g}/\text{mL}$) and with artificial sea water (3 mg in 5 ml artificial sea water) to make a 9 mL solution. After 24 hours at room temperature, the nauplii survivors can be counted macroscopically and the percent deaths at each dose and control were determined. The percentage of death was counted based on Abbott's formula (Abbot, 1925).

$$\% \text{ deaths} = [(\text{test} - \text{control}) / \text{control}] \times 100 \quad (2)$$

Statistical Analysis

The data were subjected to a one-way analysis of variance (ANOVA) and the significance of the difference between means was determined by LSD ($P < 0.05$) using SPSS version 23 (Kokoska, 2015). Values expressed are means of three replicate determinations \pm standard deviation.

RESULTS AND DISCUSSION

Proximate Composition of *C. lentillifera*

Proximate composition of FC from Natuna District of Indonesia was different than that of FC from Penghu of Taiwan except for the moisture content (**Table 1**).

Table 1. Proximate Compositions of *C. lentillifera* from Natuna District (% FC)

Proximate Composition	Percentage of FC from	
	Natuna, Indonesia	Penghu, Taiwan (Nguyen et al., 2011)
Moisture	95.01 \pm 0.170	94.28 \pm 0.24
Ash	3.41 \pm 0.160	1.27 \pm 0.02
Crude Lipid	0.79 \pm 0.002	0.09 \pm 0.01
Crude Protein	0.43 \pm 0.007	0.53 \pm 0.02
Crude fiber	14.38 \pm 3.640	0.17 \pm 0.01
Carbohydrate (exclude crude fiber)**	0.36 \pm 0.025	3.67 \pm 0.07

Values are presented as mean \pm SD (n=3). ** Calculate by difference (% carbohydrate = 100% - % moisture - % crude protein - % crude lipid - % ash - % crude fiber).

Table 2. Antimicrobial Activities of the extract DC and FC against Tested Microorganism

Microorganism Tests	MIC of Extract ($\mu\text{g}/\text{well}$)	
	DC	FC
<i>E. coli</i>	-	500
<i>S. aureus</i>	400	500
<i>B. subtilis</i>	200	50
<i>V. cholerae</i>	50	500
<i>C. freundii</i>	500	400
<i>V. harveyi</i>	500	500
<i>V. vara</i>	-	100
<i>Salmonella</i> sp.	500	-
<i>P. aeruginosa</i>	-	500
<i>A. hydrophila</i>	500	500
<i>Bacillus</i> sp.	200	-
<i>K. pneumoniae</i>	200	500
<i>C. albicans</i>	400	500

Note: -: no antimicrobial activities

FC from Natuna had generally higher levels of ash, crude lipid, and crude fiber compared with that of FC from Penghu of Taiwan. DC of Malaysia, Taiwan, and Thailand showed a generally different of their proximate composition (Nguyen et al., 2011; Ratana-arporn and Chirapart, 2006; Matanjun et al., 2009). It might be due to the different in their habitat conditions, maturity levels, and their environmental conditions (Ito & Hori, 2009).

Antimicrobial Activity

In general, extract *C. lentillifera* FC and DC showed the broad spectrum of weak antimicrobial activities against tested microorganisms (*E. coli*, *S. aureus*, *B. subtilis*, *V. cholerae*, *C. freundii*, *V. harveyi*, *P. aeruginosa*, *A. hydrophila*, *K. pneumoniae* and *C. albicans*) based on their high minimum inhibition concentration (MIC) value (Table 2). The broad spectrum of antimicrobial activities was also shown by the other extract of *Caulerpa* such as: *C. ashmeadii*, *C. paspaloides* and *C. prolifera* (Freile-pelegrin & Morales, 2004). *C. cupressoides*, *C. Mexicana*, and *C. racemose* showed their

antimicrobial activities towards *Bacillus subtilis* (Freile-pelegrin & Morales, 2004). Caulerpin and caulerpenyne were probably contributed to antimicrobial activities of *C. lentillifera* which is commonly to occur in *Caulerpa* (Paul et al., 1987).

Hemolytic Assay

The hemolytic assay can be used for toxicity studies of compounds (Situ and Bobek 2000; Djouossi et al. 2015). The advantages using of this assay is it is sensitive, cheap, quick, and easy to monitor the lysis. Both extracts of FC and DC showed its low hemolytic activities (around 5-13%) for concentrations of 100 and 150 $\mu\text{g}/\text{mL}$, but the activity increases sharply (about 96%) for both concentration of 200 $\mu\text{g}/\text{mL}$ (Table 3). The hemolytic activity will probably damage the red cell of the membrane and induces the hemolytic anemia (Zohra and Fawzia 2014). This activity probably results from the astringent phenolic content of *C. lentillifera* (Nguyen et al. 2011; Singh & Kaur 2008). According to this result, *C. lentillifera* is not recommended to be consumed in excessive amount.

Table 3. Hemolytic activity of the extract *C. lentillifera*

Sample	Percentage of Hemolysis for Concentration ($\mu\text{g}/\text{mL}$)		
	100	150	200
Extract FC	13.32 \pm 0.23a	13.48 \pm 0.27a	96.71 \pm 1.40b
Extract DC	5.11 \pm 0.56a	9.20 \pm 1.43b	96.22 \pm 1.77c

Values are presented as mean \pm SD (n=3). Values with different superscript in each column are significantly different from one another ($p < 0.05$)

Table 4. Antioxidant Activity of *C. lentillifera* Extracts Using Linoleic Acid Emulsion System

Extract	% of antioxidant activity
DC	54.23±2.28% ^a
FC	79.09±0.78% ^b
Positive Control (Vitamin C)	81.78±0.71% ^c

Values are presented as mean±SD (n=3). Values with different superscript in each row are significantly different from one another (p < 0.05)

Brine Shrimp Lethality Test (BSLT)

BSLT is a cheap and simple bioassay system which is used to detect a bioactive compound such as a cytotoxic drug (Meyer et al., 1982). BSLT result of both extracts of *C. lentillifera* showed that was only extracted of DC caused death on larvae (nauplii) of *A. salina* but no extract of FC. The extract of DC gave LC₅₀ value 258,360 µg/mL which was very low toxicity due to an LC₅₀ value greater than 100 µg/mL (Mbwambo, Moshi, Masimba, Kapingu, & Nondo, 2007). As a comparison, standard cytotoxic of drug (cyclophosphamide) and of gallic acid have LC₅₀ of 531.0 and 323.6 µg/mL, respectively (Sonibare, 2017). According to Meyer et al., the LC₅₀ value of extracts ≤ 1000 µg/mL was predicted containing bioactive compounds (Meyer et al., 1982). Caulerpenyne, a cytotoxic sesquiterpenoid, is likely responsible for this activity due to its most abundant found in the genus of *Caulerpa* (Dumay et al., 2002).

Antioxidant Activity

Antioxidant activity of an extract of *C. lentillifera* from Taiwan have been analyzed using various assays which are DPPH (1,1-diphenyl-2-picryl hydrazyl) radical scavenging, ferric reducing, hydrogen peroxide scavenging and ferrous ion chelating (FIC) activity. The extract of *C. lentillifera* showed its low antioxidant activity compared with that of a positive control for assays using DPPH radical scavenging, ferric reducing, and hydrogen peroxide scavenging (Nguyen et al., 2011). The best antioxidant activity of the extract was resulted in using FIC assay.

In this study, the extract of *C. lentillifera* was also tested for its antioxidant activity using another method namely linoleic acid emulsion system. This system measures its inhibition of lipid peroxidation by certain compounds having antioxidant activity (Loganayaki et al., 2013). Linoleic acid is an unsaturated fatty acid with 2 double bonds which is oxidized easily to be peroxides. The peroxides are able to oxidize from Fe²⁺ ions to

Fe³⁺ ions. Fe³⁺ ions then react with ion thiocyanate to produce ferri thiocyanate complex (Fe(SCN)₆) with red color. The red color intensity showed the active peroxide formed and was measured at an absorbance of 490 nm wavelength.

Both extracts showed antioxidant activity, even though the antioxidant activity value of both was significantly lower (p<0.05) than that of in the positive control, vitamin C (Table 4). According to Matanjun *et. al.*, *C. lentillifera* also contains a low level of vitamin C and vitamin E which probably contributes to antioxidant activity in this assay (Matanjun et al. 2009; Nguyen et al. 2011).

CONCLUSION

The proximate composition of *C. lentillifera* of Indonesia was different from that of Taiwan. Ash, lipid, and fiber contents of *C. lentillifera* of Indonesia were higher than that of *C. lentillifera* of Taiwan. Furthermore, *C. lentillifera* of Indonesia also contained secondary metabolites having various biological activities, even though it is not recommended to be consumed in excessive amount. According to this study, *C. lentillifera* is highly recommended to be a functional food and to be a potent secondary metabolite source with various biological activities.

REFERENCES

- Abbot, W. (1925). A Method of Computing The Effectiveness of an Insecticide. *Journal of Economic Entomology*, 18(2), 265–267.
- Bhuiyan, M. K. A., Qureshi, S., Kamal, A. H. M., Aftab, U. S., & Siddique, M. A. M. (2016). Proximate Chemical Composition of Sea Grapes *Caulerpa racemosa* (J. Agardh, 1873) Collected from a Sub-Tropical Coast. *Virology and Mycology*, 5(2). <https://doi.org/10.4172/2161-0517.1000158>
- Bonev, B., Hooper, J., & Parisot, J. (2008).

- Principles of Assessing Bacterial Susceptibility to Antibiotics using the Agar Diffusion Method. *Journal of Antimicrobial Chemotherapy*, 61(March), 1295–1301.
<https://doi.org/10.1093/jac/dkn090>
- Daud, D., Fazuliana, N. U. R., Arsad, M., Ismail, A., & Tawang, A. (2016). Antipyretic Action of *Caulerpa lentillifera*, *Hibiscus rosasinensis*, and *Piper sarmentosum* Aqueous Extract in Mice. *Asian Journal of Pharmaceutical and Clinical Research*, 9(1), 9–11.
- Djouossi, M. G., Tamokou, J., Ngnokam, D., Kuate, J., Tapondjou, L. A., Harakat, D., & Voutquenne-nazabadioko, L. (2015). Antimicrobial and Antioxidant Flavonoids from the Leaves of *Oncoba spinosa* Forsk. (Salicaceae). *BMC Complementary and Alternative Medicine*, 15, 4–11.
<https://doi.org/10.1186/s12906-015-0660-1>
- Doty, S. M. (1966). Caulerpicin, a Toxic Constituent of *Caulerpa*. *Nature*, 211, 990.
- Doty, S. M., & Aguilar-santos, G. (1970). Transfer of Toxic Alga Substances in Marine Food Chains. *Pacific Science*, 24, 351–355.
- Dumay, O., Erard, G., Pergent-martini, C., & Amade, P. (2002). Variations in Caulerpenyne Contents in *Caulerpa taxifolia* and *Caulerpa racemosa*. *Journal of Chemical Ecology*, 28(2), 343–352.
- Freile-pelegrin, Y., & Morales, J. L. (2004). Antibacterial Activity in Marine Algae from the Coast of Yucatan, Mexico. *Botanica Marina*, 47, 140–146.
<https://doi.org/10.1515/BOT.2004.014>
- Ito, K., & Hori, K. (2009). Seaweed : Chemical Composition and Potential Food Uses. *Food Reviews International*, 5(1), 101–144.
- Kılınç, B., Cirik, S., & Turan, G. (2013). Seaweeds for Food and Industrial Applications. In *Food Industry* (pp. 735–748). INTECH.
- Kokoska, S. (2015). *Introductory Statistics: A Problem-Solving Approach* (Second). New York: W.H Freeman and Company.
- Loganayaki, N., Siddhuraju, P., & Manian, S. (2013). Antioxidant Activity and Free Radical Scavenging Capacity of Phenolic Extracts from *Helicteres isora* L. and *Ceiba pentandra* L. *Journal of Food Science and Technology*, 50(4), 687–695.
<https://doi.org/10.1007/s13197-011-0389-x>
- Matanjun, P., Mohamed, S., Mustapha, N. M., & Muhammad Kharidah. (2009). Nutrient Content of Tropical Edible Seaweeds, *Eucheuma cottonii*, *Caulerpa lentillifera* and *Sargassum polycystum*. *Journal of Applied Phycology*, 21, 75–80.
<https://doi.org/10.1007/s10811-008-9326-4>
- Mbwambo, Z. H., Moshi, M. J., Masimba, P. J., Kapingu, M. C., & Nondo, R. S. O. (2007). Antimicrobial Activity and Brine Shrimp Toxicity of Extracts of *Terminalia brownii* roots and stem. *BMC Complementary and Alternative Medicine*, 7(9), 5–9.
<https://doi.org/10.1186/1472-6882-7-9>
- Meyer, B. N., Ferrigni, N. A., Putnam, J. E., Jacobsen, L. B., Nichols, D. E., & McLaughlin, J. L. (1982). Brine Shrimp : A Convenient General Bioassay for Active Plant Constituents. *Journal of Medicinal Plant Research*, 45, 31–34.
- Mohamed, S., Hashim, S. N., & Rahman, A. (2012). Seaweeds : A Sustainable Functional Food for Complementary and Alternative Therapy. *Trends in Food Science & Technology*, 23(2), 83–96.
<https://doi.org/10.1016/j.tifs.2011.09.001>
- Nguyen, T. Van, Ueng, J.-P., & Tsai, G.-J. (2011). Proximate Composition, Total Phenolic Content, and Antioxidant Activity of Sea grape (*Caulerpa lentillifera*). *Journal of Food Science*, 76 (November), C950-c958.
<https://doi.org/10.1111/j.1750-3841.2011.02289.x>
- Okeke, M. ., Iroegbu, C. ., Eze, E. ., Okoli, A. ., & Esimone, C., (2001). Evaluation of Extracts of Root of *Landonphin owerrience* for Antibacterial Activity. *Journal of Ethnopharmacology*, 78, 119–127. [https://doi.org/10.1016/S0378-8741\(01\)00307-5](https://doi.org/10.1016/S0378-8741(01)00307-5)
- Paul, J. V., Littler, M. M., Littler, S. D., & Fenical, W. (1987). Evidence for Chemical Defense in Tropical Green Alga *Caulerpa ashmeadii* (Caulerpaceae: Chlorophyta): Isolation of New Bioactive Sesquiterpenoids. *Journal of Chemical Ecology*, 13(5), 1171–1185.

- Ratana-arporn, P., & Chirapart, A. (2006). Nutritional Evaluation of Tropical Green Seaweeds *Caulerpa lentillifera* and *Ulva reticulata*. *Kasetsart J. (Nat. Sci)*, 40(September), 75–83.
- Saengkhae, C., Arunnopparat, W., & Sungkhajorn, P. (2007). Antioxidative Activity of the Leaf of *Nelumbo nucifera* Gaertn. on Oxidative Stress-Induced Erythrocyte Hemolysis in Hypertensive and Normotensive Rats. *Thai Journal of Physiological Sciences*, 20(2), 70–78.
- Singh, R. P., & Kaur, G. (2008). Hemolytic Activity of Aqueous Extract of *Livistona chinensis* Fruits. *Food and Chemical Toxicology*, 46, 553–556. <https://doi.org/10.1016/j.fct.2007.08.037>
- Situ, H., & Bobek, A. L. (2000). In Vitro Assessment of Antifungal Therapeutic Potential of Salivary Histatin-5, Two Variants of Histatin-5, and Salivary Mucin (MUC7) Domain 1. *Antimicrobial Agents and Chemotherapy*, 44(6), 1485–1493.
- Sonibare, A. A. A. M. A. (2017). In Vitro Antioxidant Activity, Brine Shrimp Lethality and Assessment of Bioactive Constituents of Three wild Dioscorea Species. *Journal of Food Measurement and Characterization*, 11(2), 685–695. <https://doi.org/10.1007/s11694-016-9438-5>
- Zohra, M., & Fawzia, A. (2014). Hemolytic Activity of Different Herbal Extracts used in Algeria. *International Journal of Pharma Science and Research*, 5(08), 495–500.